

Breastfeeding: a natural defence against obesity?

Gabriella D'Angelo, Sara Manti, Andrea Barbalace, Ignazio Barberi

Neonatal and Pediatric Intensive Care Unit, Department of Pediatrics, University of Messina, Italy

Abstract

Today, obesity represents one of the most serious health problems facing both children and adults. Childhood obesity has several causes, including genetic factors, dietary habits, personal behaviours, and interaction of all of these. It often leads to adult obesity, which causes health problems including heart disease, diabetes, and even early death. Thus, many studies have investigated possible measures to prevent childhood obesity, and breastfeeding is considered an important early preventive intervention. Despite the fact that several milk formulas have been demonstrated to be safe and effective for feeding both term and premature infants, for its immunological and nutritional qualitative advantages, human milk is nowadays universally recognized as the optimal feeding choice for healthy, sick and preterm infants.

To date, it is however still unclear whether breastfeeding can prevent childhood obesity. In fact, literature data provide controversial results, probably due to several confounding factors, including maternal habits, age, level of education, lifestyle, race, parity, pregnancy complications, types of delivery, and infant health factors.

Thus, whether breastfeeding protects against obesity is still unclear. Further researches, by reducing the influence of confounding factors and improving the accuracy of the effect estimate, are needed to confirm the validity of the role of breastfeeding in reducing the risk of developing childhood overweight.

This review briefly summarizes what is known on the possible relationship between breastfeeding and prevention of obesity development.

Keywords

Breastfeeding, human milk, overweight, obesity, adipokines, children.

Corresponding author

Sara Manti, Neonatal and Pediatric Intensive Care Unit, Department of Pediatrics, University of Messina, Italy; email: saramanti@hotmail.it.

How to cite

D'Angelo G, Manti S, Barbalace A, Barberi I. Breastfeeding: a natural defence against obesity? *J Pediatr Neonat Individual Med.* 2016;5(1):e050106. doi: 10.7363/050106.

Introduction

Childhood obesity is a global epidemic health problem and rising trends in overweight is apparent in both developed and developing countries [1]. Many strategies addressing prevention of childhood obesity and related risk of chronic diseases have been proposed, including breastfeeding, which is widely acknowledged as the ideal form of nutrition for infants [1].

Despite the fact that several milk formulas have been demonstrated to be safe and effective for feeding both term and premature infants [2], for its immunological and nutritional qualitative advantages, human milk (HM) is nowadays universally recognized as the optimal feeding choice for healthy, sick and preterm infants. Breast milk is uniquely suited to the human infant, and, as a dynamic, bioactive fluid, it changes in composition from colostrum to late lactation, and varies within feeds, diurnally, and between mothers [3]. It also confers neural, psychological, and developmental benefits [4].

Furthermore, breastfeeding has been shown to be protective against several illnesses, including upper and lower respiratory infections, atopic (asthma and eczema) and immunological (celiac disease) disorders, as well as overweight/obesity [3]. In particular, it has been reported that, although breastfed infants grow more rapidly during four months of life, at age 1 they gain weight less than formula fed babies [5, 6].

In light of these and similar results, it has been assumed that breastfeeding can prevent overweight. However, as obesity is considered a multifactorial disorder [7], whether breastfeeding can help to prevent obesity childhood is still unclear [8]. In fact, data from literature concerning a possible correlation between breastfeeding and overweight/obesity prevention provided controversial results, probably due to several confounding causes. These include: maternal body mass index (BMI), age, race/ethnicity, lifestyle, level of education, smoking, sedentary or physically activity, parity, types of delivery, pregnancy complications, and infant health factors [9].

Possible effects of breastfeeding on weight gain regulation

Several hypothesis have been proposed to explain the protective role of breastfeeding against obesity.

Firstly, protein intake as well as energy metabolism are lower among breastfed subjects [10, 11]. Despite the fact that breast milk provides a moderate amount of calories and nutrients for the infant (e.g., protein, sugar, fat, and water) and also changes with time and mother's diet [3], the content of protein in formula feeding is approximately 1.6 times higher than baby's needs [12]. In particular, in longitudinal studies, a significant relationship was found between dietary protein intake at the age of 10 months and later BMI and distribution of body fat [13]. This confirms that a high intake of protein in early childhood might increase the risk of obesity in later life.

Secondly, studies revealed the presence, in human milk, of hormones (e.g., leptin, adiponectin, resistin, ghrelin) which, through early control of satiety, have a positive effect on the programming of energy balance regulation in childhood and adulthood, thereby protecting against later obesity [14].

In particular, leptin, an adipocyte-derived hormone also present in human milk, passes from breast-milk to infant blood and, by acting as a satiety signal, exerts both short- and long-term effects on energy balance and body weight regulation [15, 16]. The existence of leptin in breast milk might have a significant role in growth, food intake and regulation of satiety in infants during the early lactation period [17, 18].

Adiponectin, the most abundant adipose-specific protein, circulates in very high concentrations in human milk and levels are positively associated with insulin sensitivity and inversely related to the degree of adiposity [19]. An inverse relationship between adiponectin levels in milk and adiposity of the breastfed infant has been observed; this might be due to modulation of infant metabolism by milk adiponectin, and may be related to the observed protection against obesity from breastfeeding [20].

Resistin, a cytokine synthesized by adipocytes, correlates positively with hormone status (estradiol, progesterone, prolactin, thyroxine, triiodothyronine, cortisol, and leptin); thus, it may be involved in appetite regulation and in the metabolic balance of newborns [21, 22]. It has been reported that resistin has a significant role in

energy and glucose homeostasis, since it impairs the activation of AMP-activated protein kinase (AMPK) which is a key regulator of glucose uptake and skeletal muscle fatty acid metabolism [23]. Recently, a positive correlation between serum resistin level of breastfed infants and breast milk resistin levels has been founded [24].

Another component of human milk is ghrelin which, acting as an appetite-stimulatory signal and playing a role in short- and long-term regulation of food intake, influences the energy balance [25]. Ghrelin would appear to be involved in postnatal growth, and a positive correlation between ghrelin level in breast milk and infant weight gain has been observed [26]. However, serum ghrelin levels in formula fed infants is significantly higher compared to breastfed infants [27]. This finding might suggest that formula fed infants, who receive a higher amount of ghrelin, have a greater feeding stimulus than breastfed infants and a consequent increase in weight and growth rate.

Moreover, human milk, through both lipid and not lipid molecules, including epidermal growth factor and tumour necrosis factor α , increases the growth factors synthesis which, in turns, inhibits preadipocyte differentiation [28].

Nevertheless, several other peptide/protein hormones have recently been identified in human breast milk, including obestatin, nesfatin, irisin, adropin, copeptin, pituitary adenylate cyclase-activating polypeptide, apelin, motilin and cholecystokinin, that may be involved in the mechanism of weight gain regulation [29].

It is known that breastfeeding can influence the prevalence of both underweight and overweight status in later life [30, 31]. For example, a major discharge of growth hormone, insulin-like growth factor 1 and insulin are present in the bloodstream of breastfed newborns, promoting a major fat deposition and an increased number of adipocytes [32].

In addition to biological mechanisms, breastfeeding also influences behavioural and physiological feeding patterns. Breastfed infants tend to better adjust the amount of human milk and calories they ingest, in comparison to formula fed infants, who are often obligate to finish a bottle even if satiated.

Finally, it has been reported that breastfeeding affects timing of solid food intake. Exclusive breastfeeding precludes inappropriate complementary feeding practices such as early introduction of complementary foods that could

lead to unhealthy weight gain [33]. Huh and colleagues found that early introduction of solids was associated with higher mean BMI at 3 years of age among formula fed infants, but not breastfed infants [34].

Moreover, it has been proposed that differences in dietary preferences between breast and formula fed infants could also justify breastfeeding-risk obesity association. Scholtens et al. [35] reported that, at 7 years of age, breastfed children for more than 16 weeks had a major intake of fruit and vegetables than formula fed newborns.

Breastfeeding and gain weight: state of art

Evidence on breastfeeding and obesity are, to date, contradictory, probably due to several confounding variables which can influence body weight in children and contribute to overweight/obesity prevention. In 2004 and 2005, two meta-analyses reported that breastfeeding could help to prevent obesity childhood [36, 37]. Thereafter, through the meta-analyses of childhood risk factors for obesity, Wheng et al. [38] founded that breastfeeding infants had a 15% decrease in the odds of childhood overweight in comparison to bottle fed babies. In this paper, ten studies [30, 39-47], comparing the effects of breastfeeding and other types of nutrition during the first 12 months of life, were particularly analyzed. From the collected data the protective role of breastfeeding against overweight in childhood resulted unclear. In fact, 5 studies [30, 41, 44, 45, 47] reported a beneficial effect of breastfeeding against obesity, while the other 5 did not confirm the beneficial role [39, 40, 42, 43, 46]. Among the studies [39, 44, 48-50] which analyzed the possible influence of breastfeeding duration on childhood overweight, only one [39] reported a significant decrease in the OR of overweight at 2 years of age if infants who were breastfed for more than 6 months were compared with those breastfed for less than 3 months. With the same purpose, Yan et al. [51] conducted a meta-analysis, including 25 studies (10 studies cross-sectional surveys and 15 cohort studies) with a total of 226,508 participants. They investigated the association between breastfeeding duration (categorized as < 3 months, 3-5 months, 5-7 months and \geq 7 months) and obesity. In detail, authors assessed that the risk of childhood obesity was lower in breastfed children by 22% compared with those who were never breastfed. Moreover, infants who were breastfed for more than 7 months

had significantly less likelihood to develop obesity, while in those breastfed for less than 3 months the obesity risk decreased of 10%. Authors conclude that breastfeeding is a significant protective factor against obesity in children, indicating also a breastfeeding dose-response effect.

Conversely, in a recent national survey enrolling 14,880 children, Djalalinia et al. [52] assessed that duration of breastfeeding, categorized into five groups of 0 months, < 6 months, 6-12 months, 12-18 months, and 18-24 months, had no significant association with anthropometric measures such as height, weight, blood pressure, and abdominal obesity. Compared to previous studies [39, 51], the authors also hypothesized that the controversies regarding the relationship between anthropometric measures and breastfeeding might result from the presence of several parameters, including ethnicity, age, maternal glucose tolerance, hormonal and body composition changes, acting as a confounding factors. Confirming this hypothesis, Assunção et al. [53] reported that, after adjusting for potential confounding parameters (e.g., anthropometric, socioeconomic, demographic, behavioural, and health-related factors), children, exposed to exclusive breastfeeding for more than 6 months, showed a lower prevalence of overweight in the second year of life.

In addition, several studies assessed that exclusive breastfeeding appears to have a stronger protective effect than breastfeeding combined with formula feeding [37, 45, 54, 55]. However, these results were not confirmed by the findings of several studies, in which the obesity risk was not found decreased in children breastfed for less than 4 months [49], for 4 months [56], for 6 months [50], for less than 12 months or major than 12 months [57] of age.

In conclusion, while the beneficial effects of breastfeeding children are well documented in several diseases [3], data that breastfeeding, also in a dose-dependent manner, might decrease risk of both short- and long-term obesity are controversial, showing both protective and null effects. These discrepancies may be related to the diverse enrolled populations (with unique genetic and environmental backgrounds), different sample sizes and absence/presence of confounding factors (e.g., maternal education, maternal age, maternal pre- and post-partum weight, parity, ethnicity, and delivery type). Further research, by reducing the influence of confounding factors and improving the accuracy

of the effect estimate, are needed to confirm the validity of the role of breastfeeding in reducing the risk of developing childhood overweight. In the mean time, breastfeeding, despite the rare results of its uncorrect management, including hypernatremic dehydration [58-60], still represents the most ancient and natural mode of feeding for the newborn, and should always be encouraged [61]. Moreover, the results of the abovementioned studies seem also to confirm that breastfeeding is not only nutritionally sound but economically beneficial as well [62], in particular, by reducing the sanitary costs related to obesity. On this regard, international data collected by the International Association for the Study of Obesity show that 36% of boys and 34% of girls are overweight or obese in Italy, compared with 23% of boys and 21% of girls, on average, in other countries [63]. This critical incidence causes an obesity cost equal to 2,500,000,000 euros/year. Nevertheless several programmes have been proposed to cut health expenditures, their health effects take a longer time. In fact, while the treatment for obesity is time-consuming and inconclusive, the prevention of obesity is the most effective solution [64]. Therefore, breastfeeding, through protective effect against obesity, could be also represent an effective prevention strategy.

Declaration of interest

The Authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article. The Authors declared that they did not receive any financial support.

References

1. World Health Organization. Obesity: preventing and managing the global epidemic: report of a WHO consultation. World Health Organ Tech Rep Ser. 2000;894:i-253.
2. Marseglia L, Pagano G, Arco A, Barberi I, Biasucci G, Riboni S, Mondello I, Fiamingo C, Moro G. A new formula for premature infants: effects on growth and nutritional status. *J Matern Fetal Neonatal Med.* 2014 Sep 22. [Epub ahead of print].
3. Horta BL, Victora CG. Long-term effects of breastfeeding: a systematic review. Geneva: World Health Organization, 2013.
4. Mortensen EL, Michaelsen KF, Sanders SA, Reinisch JM. The association between duration of breastfeeding and adult intelligence. *JAMA.* 2002;287:2365-71.
5. Dewey KG. Growth characteristics of breastfed compared to formula fed infants. *Biol Neonate.* 1998;74:94-105.
6. Duncan B, Schaefer C, Sibley B, Fonseca NM. Reduced growth velocity in exclusively breastfed infants. *Am J Dis Child.* 1984;138:309-13.

7. Marseglia L, Manti S, D'Angelo G, Nicotera A, Parisi E, Di Rosa G, Gitto E, Arrigo T. Oxidative stress in obesity: a critical component in human diseases. *Int J Mol Sci*. 2014;16(1):378-400.
8. Marseglia L, Manti S, D'Angelo G, Cuppari C, Salpietro V, Filippelli M, Trovato A, Gitto E, Salpietro C, Arrigo T. Obesity and breastfeeding: The strength of association. *Women Birth*. 2015 Jan 13. [Epub ahead of print].
9. Monasta L, Batty GD, Cattaneo A, Lutje V, Ronfani L, Van Lenthe FJ. Early-life determinants of overweight and obesity: a review of systematic reviews. *Obes Rev*. 2010;11:695-708.
10. Whitehead RG. For how long is exclusive breast-feeding adequate to satisfy the dietary energy needs of the average young baby? *Pediatr Res*. 1995;37:239-43.
11. Rolland-Cachera MF, Deheeger M, Akrouit M, Bellisle F. Influence of macronutrients on adiposity development: a follow up study of nutrition and growth from 10 months to 8 years of age. *Int J Obes Relat Metab Disord*. 1995;19:573-8.
12. Agostoni C, Braegger C, Decsi T, Kolacek S, Koletzko B, Michaelsen KF, Mihatsch W, Moreno LA, Puntis J, Shamir R, Szajewska H, Turck D, van Goudoever J. Breast-feeding: A Commentary by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr*. 2009;49:112-25.
13. Deheeger M, Akrouit M, Bellisle F, Rossignol C, Rolland-Cachera MF. Individual patterns of food intake development in children: a 10 months to 8 years of age follow-up study of nutrition and growth. *Physiol Behav*. 1996;59:403-7.
14. Savino F, Liguori SA. Update on breast milk hormones: leptin, ghrelin and adiponectin. *Clin Nutr*. 2008;27(1):42-7.
15. Ilcol YO, Hizli ZB, Ozkan T. Leptin concentration in breast milk and its relationship to duration of lactation and hormonal status. *Int Breastfeed J*. 2006;1:21.
16. Mamì C, Manganaro R, Saitta G, Marseglia L, Martino F, Gemelli M. Plasma leptin, insulin, and neuropeptide Y concentrations in infants. *Arch Dis Child Fetal Neonatal Ed*. 2005;90(1):F86-7.
17. Savino F, Costamagna M, Prino A, Oggero R, Silvestro L. Leptin levels in breastfed infants. *Acta Paediatr*. 2002;91:897-902.
18. Mamì C, Manganaro R, Marseglia L, Saitta G, Gemelli M, Martino F. Plasma leptin, insulin, and neuropeptide Y response to feeding in newborn infants. *Arch Dis Child Fetal Neonatal Ed*. 2006;91(5):F389.
19. Martin LJ, Woo JG, Geraghty SR, Altaye M, Davidson BS, Banach W, Dolan LM, Ruiz-Palacios GM, Morrow AL. Adiponectin is present in human milk and is associated with maternal factors. *Am J Clin Nutr*. 2006;83(5):1106-11.
20. Newburg DS, Woo JG, Morrow AL. Characteristics and potential functions of human milk adiponectin. *J Pediatr*. 2010;156(Suppl): 41-6.
21. Ilcol YO, Hizli ZB, Eroz E. Resistin is present in human breast milk and it correlates with maternal hormonal status and serum level of C-reactive protein. *Clin Chem Lab Med*. 2008;46(1):118-24.
22. Mamì C, Marseglia L, Manganaro R, Saitta G, Martino F, Gargano R, Gemelli M. Serum levels of resistin and its correlation with adiponectin and insulin in healthy full term neonates. *Early Hum Dev*. 2009;85(1):37-40.
23. Dick DJ, Heigenhauser GJF, Bruce CR. The role of adipokines as regulators of skeletal muscle fatty acid metabolism and insulin sensitivity. *Acta Physiol*. 2006;186:5-16.
24. Savino F, Sorrenti M, Benetti S, Lupica MM, Liguori SA, Oggero R. Resistin and leptin in breast milk and infants in early life. *Early Hum Dev*. 2012;88:779-82.
25. Cummings DE. Ghrelin and the short- and long-term regulation of appetite and body weight. *Physiol Behav*. 2006;89(1):71-84.
26. Cesur G, Ozguner F, Yilmaz N, Dundar B. The relationship between ghrelin and adiponectin levels in breast milk and infant serum and growth of infants during early postnatal life. *J Physiol Sci*. 2012;62:185-90.
27. Savino F, Petrucci E, Lupica MM, Nanni GE, Oggero R. Assay of ghrelin concentration in infant formulas and breast milk. *World J Gastroenterol*. 2011;17:1971-5.
28. Hauner H, Rohrig K, Petruschke T. Effects of epidermal growth factor (EGF), platelet-derived growth factor (PDGF) and fibroblast growth factor (FGF) on human adipocyte development and function. *Eur J Clin Invest*. 1995;25:90-6.
29. Çatlı G, Olgaç Dünder N, Dünder BN. Adipokines in breast milk: an update. *J Clin Res Pediatr Endocrinol*. 2014;6(4):192-201.
30. Grummer-Strawn LM, Mei Z. Does breastfeeding protect against pediatric overweight? Analysis of longitudinal data from the Centers for Disease Control and Prevention Pediatric Nutrition Surveillance System. *Pediatrics*. 2004;113:e81-6.
31. Beyerlein A, Toschke AM, von Kries R. Breastfeeding and childhood obesity: shift of the entire BMI distribution or only the upper parts? *Obesity (Silver Spring)*. 2008;16:2730-3.
32. Lucas A, Sarson DL, Blackburn AM, Adrian TE, Aynsley-Green A, Bloom SR. Breast vs bottle: endocrine responses are different with formula feeding. *Lancet*. 1980;1:1267-9.
33. Moreno MA, Furtner F, Rivara FP. Breastfeeding as Obesity Prevention. *Arch Pediatr Adolesc Med*. 2011;165(8):772.
34. Huh SY, Rifas-Shiman SL, Taveras EM, Oken E, Gillman MW. Timing of solid food introduction and risk of obesity in preschool-aged children. *Pediatrics*. 2011;127(3):e544-51.
35. Scholtens S, Brunekreef B, Smit HA, Gast GC, Hoekstra MO, de Jongste JC, Postma DS, Gerritsen J, Seidell JC, Wijga AH. Do differences in childhood diet explain the reduced overweight risk in breastfed children? *Obesity (Silver Spring)*. 2008;16:2498-503.
36. Arenz S, Ruckerl R, Koletzko B, Von Kries R. Breastfeeding and childhood obesity – a systematic review. *Int J Obes (Lond)*. 2004;28:1247-56.
37. Owen CG, Martin RM, Whincup PH, Smith GD, Cook DG. Effect of infant feeding on the risk of obesity across the life course: a quantitative review of published evidence. *Pediatrics*. 2005;115:1367-77.
38. Weng SF, Redsell SA, Swift JA, Yang M, Glazebrook CP. Systematic review and meta-analyses of risk factors for childhood overweight identifiable during infancy. *Arch Dis Child*. 2012;97(12):1019-26.
39. Weyermann M, Rothenbacher D, Brenner H. Duration of breastfeeding and risk of overweight in childhood: a prospective birth cohort study from Germany. *Int J Obes*. 2006;30:1281-7.

40. Shields L, O'Callaghan M, Williams GM, Najman JM, Bor W. Breastfeeding and obesity at 14 years: a cohort study. *J Paediatr Child Health*. 2006;42:289-96.
41. Hawkins SS, Cole TJ, Law C; Millennium Cohort Study Child Health Group. An ecological systems approach to examining risk factors for early childhood overweight: findings from the UK Millennium Cohort Study. *J Epidemiol Community Health*. 2009;63:147-55.
42. Kwok MK, Schooling CM, Lam TH, Leung GM. Does breastfeeding protect against childhood overweight? Hong Kong's 'Children of 1997' birth cohort. *Int J Epidemiol*. 2010;39:297-305.
43. Burke V, Beilin LJ, Simmer K, Oddy WH, Blake KV, Doherty D, Kendall GE, Newnham JP, Landau LI, Stanley FJ. Breastfeeding and overweight: longitudinal analysis in an Australian birth cohort. *J Pediatr*. 2005;147:56-61.
44. Taveras EM, Rifas-Shiman SL, Scanlon KS, Grummer-Strawn LM, Sherry B, Gillman MW. To what extent is the protective effect of breastfeeding on future overweight explained by decreased maternal feeding restriction? *Pediatrics*. 2006;118:2341-8.
45. Armstrong J, Reilly JJ. Breastfeeding and lowering the risk of childhood obesity. *Lancet*. 2002;359:2003-4.
46. Reilly JJ, Armstrong J, Dorosty AR, Emmett PM, Ness A, Rogers I, Steer C, Sherriff A; Avon Longitudinal Study of Parents and Children Study Team. Early life risk factors for obesity in childhood: cohort study. *Br Med J (Clinical Research Edition)*. 2005;330:1357-9.
47. Bergmann KE, Bergmann RL, Von Kries R, Böhm O, Richter R, Dudenhausen JW, Wahn U. Early determinants of childhood overweight and adiposity in a birth cohort study: role of breastfeeding. *Behav Modif*. 2003;27:54-67.
48. Buyken AE, Karaolis-Danckert N, Remer T, Bolzenius K, Landsberg B, Kroke A. Effects of breastfeeding on trajectories of body fat and BMI throughout childhood. *Obesity*. 2008;16:389-95.
49. Huus K, Ludvigsson JF, Enskär K, Ludvigsson J. Exclusive breastfeeding of Swedish children and its possible influence on the development of obesity: a prospective cohort study. *BMC Pediatrics*. 2008;8:42.
50. Toschke AM, Martin RM, von Kries R, Wells J, Smith GD, Ness AR. Infant feeding method and obesity: body mass index and dual-energy X-ray absorptiometry measurements at 9–10 y of age from the Avon Longitudinal Study of Parents and Children (ALSPAC). *Am J Clin Nutr*. 2007;5:1578-85.
51. Yan J, Liu L, Zhu Y, Huang G, Wang PP. The association between breastfeeding and childhood obesity: a meta-analysis. *BMC Public Health*. 2014;14:1267.
52. Djalalinia S, Qorbani M, Heshmat R, Motlagh ME, Ardalan G, Bazyar N, Taheri M, Asayesh H, Kelishadi R. Association of Breast Feeding and Birth Weight with Anthropometric Measures and Blood Pressure in Children and Adolescents: The CASPIAN-IV Study. *Pediatr Neonatol*. 2015 Mar 13. [Epub ahead of print].
53. Assunção ML, Ferreira HS, Coutinho SB, Santos LM, Horta BL. Protective effect of breastfeeding against overweight can be detected as early as the second year of life: a study of children from one of the most socially-deprived areas of Brazil. *J Health Popul Nutr*. 2015;33(1):85-91.
54. Hediger ML, Overpeck MD, Kuczumarski RJ, Ruan WJ. Association between infant breastfeeding and overweight in young children. *J Am Med Assoc*. 2001;285(19):2453-60.
55. Liese AD, Hirsch T, von Mutius E, Keil U, Leupold W, Weiland SK. Inverse association of overweight and breast feeding in 9 to10-year old children in Germany. *Int J Obes Relat Metab Disord*. 2001;25:1644-50.
56. Burdette HL, Whitaker RC, Hall WC, Daniels SR. Breastfeeding, introduction of complementary foods, and adiposity at 5 y of age. *Am J Clin Nutr*. 2006;83:550-8.
57. Gillman MW, Rifas-Shiman SL, Kleinman K, Oken E, Rich-Edwards JW, Taveras EM. Developmental origins of childhood overweight: potential public health impact. *Obesity*. 2008;16:1651-6.
58. Manganaro R, Mami C, Marrone T, Marseglia L, Gemelli M. Incidence of dehydration and hypernatremia in exclusively breastfed infants. *J Pediatr*. 2001;139(5):673-5.
59. Manganaro R, Marseglia L, Mami C, Palmara A, Paolata A, Loddo S, Gargano R, Mondello M, Gemelli M. Breast milk sodium concentration, sodium intake and weight loss in breastfeeding newborn infants. *Br J Nutr*. 2007;97(2):344-8.
60. Manganaro R, Marseglia L, Mami C, Saitta G, Gargano R, Gemelli M. Serum alpha-fetoprotein (AFP) levels in breastfed infants with prolonged indirect hyperbilirubinemia. *Early Hum Dev*. 2008;84(7):487-90.
61. Manganaro R, Marseglia L, Mami C, Paolata A, Gargano R, Mondello M, Puliafito A, Gemelli M. Effects of hospital policies and practices on initiation and duration of breastfeeding. *Child Care Health Dev*. 2009;35(1):106-11.
62. Yan J, Liu L, Zhu Y, Huang G, Wang PP. The association between breastfeeding and childhood obesity: a meta-analysis. *BMC Public Health*. 2014;14:1267.
63. OECD. Health at a Glance 2013: OECD Indicators. Paris: OECD Publishing, 2013.
64. Cole TJ. Early causes of child obesity and implications for prevention. *Acta Paediatr Suppl*. 2007;96:2-4.